

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/103627/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Muller, I. ORCID: <https://orcid.org/0000-0003-2926-0722>, Kilburn, L. S., Taylor, P. N., Barrett-Lee, P. J. ORCID: <https://orcid.org/0000-0002-6246-7357>, Bliss, J. M., Ellis, P., Ludgate, M. E. and Dayan, C. M. ORCID: <https://orcid.org/0000-0002-6557-3462> 2017. TPOAb and thyroid function are not associated with breast cancer outcome: evidence from a large-scale study using data from the Taxotere as Adjuvant Chemotherapy Trial (TACT, CRUK01/001). European Thyroid Journal 6 (4) , pp. 197-207.
10.1159/000460246 file

Publishers page: <http://dx.doi.org/10.1159/000460246>
<<http://dx.doi.org/10.1159/000460246>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



TPOAb and thyroid function are not associated with breast cancer outcome; evidence from a large-scale study using data from the Taxotere as Adjuvant Chemotherapy Trial (TACT, CRUK01/001)

Ilaria Muller¹, Lucy S. Kilburn², Peter N. Taylor¹, Peter J. Barrett-Lee³, Judith M. Bliss², Paul Ellis⁴, Marian E. Ludgate¹, Colin M. Dayan¹

¹ Thyroid Research Group, School of Medicine, Cardiff University, Cardiff, UK

² Institute of Cancer Research - Clinical Trials & Statistics Unit (ICR-CTSU), London, UK

³ Academic Breast Department, Velindre Cancer Centre, Cardiff, UK

⁴ Guy's Hospital & King's College, London, UK

Running title: Thyroid autoimmunity and breast cancer outcome

DISCLOSURE STATEMENT: The authors have nothing to disclose.

Word count: 2499

CORRESPONDING AUTHOR: Dr. Ilaria Muller, MBBS, PhD

Thyroid Research Group, Division of Infection & Immunity

School of Medicine, Cardiff University

Main building Room 256 C2 Link Corridor, University Hospital of Wales, Heath Park, CF14

4XN, Cardiff, United Kingdom (UK)

Phone: +44 (0)29 2074 5409, +44 (0)29 2074 5457

Fax: +44 (0)29 2074 4671

Email address: mulleri4@cardiff.ac.uk

27

28 **Key words:** Autoimmune thyroid disease; Autoimmunity; Hyperthyroidism; Hypothyroidism;

29 Thyroid function; Thyroid peroxidase antibodies; Breast cancer.

ABSTRACT

Background: Small-scale studies correlated the presence of thyroid autoimmunity with both improved or worsened breast cancer outcome.

Objectives: We aimed to clarify this association in a large cohort using the phase-III randomized controlled “Taxotere as Adjuvant Chemotherapy Trial” (TACT, CRUK01/001).

Methods: TACT women >18-years-old with node-positive or high risk node-negative early breast cancer (pT1-3a,pN0-1,M0), with stored plasma (n=1974), taken 15.5 [7.0-24.0] months (median [IQR]) after breast surgery were studied. Patients had also received chemotherapy (100%), radiotherapy (1745/1974 [88.4%]), hormonal therapy (1378/1974 [69.8%]), or trastuzumab (48/1974 [2.4%]). History of thyroid diseases and/or related treatments was not available.

The prognostic significance of autoantibodies to thyroid peroxidase (TPOAb; positive ≥ 6 kIU/L), free-thyroxine and thyrotropin (combined: euthyroid, hypothyroid, hyperthyroid) was evaluated for disease-free survival (DFS), overall-survival (OS), time-to-recurrence (TTR), with Cox regression models in univariate and multivariable analyses. The extended median follow-up was 97.5 months.

Results: No difference in DFS was found by TPOAb status (unadjusted-hazard ratio [HR]: 0.97, 95%CI: 0.78-1.19, P=0.75) and/or thyroid function (unadjusted-HR [hypothyroid versus euthyroid]: 1.15, 95%CI: 0.79-1.68, P=0.46; unadjusted-HR [hyperthyroid versus euthyroid]: 1.14, 95%CI: 0.82-1.61, P=0.44). Similar results were obtained for OS, TTR, multivariable analyses, when TPOAb titre by tertiles was considered and in a subgroup of 123 patients with plasma collected before adjuvant treatments.

Conclusions: No evidence for a prognostic role of TPOAb and/or thyroid function in moderate-high risk early breast cancer was found in the largest and longest observational study to date.

INTRODUCTION

An association between breast cancer (BC) and benign thyroid disorders has been debated for decades, reported in several [1,2], but not all [3] studies; the most recent meta-analyses and reviews reached contrasting conclusions [1,4-6]. Hypothyroidism was found to correlate with both an increased [7,8] or reduced [9-11] risk of developing BC, whilst other authors did not report a significant correlation [12,13]. BC has been particularly associated with thyroid autoimmunity (TA); a higher prevalence of anti-thyroid peroxidase (TPO) autoantibodies (TPOAb) was found among BC patients, compared with healthy controls [8,14]. Furthermore, a better BC outcome has been reported in TPOAb positive (TPOAb+) versus TPOAb negative (TPOAb-) patients in some [15-18], but not all [19] studies.

Currently no validated major blood prognostic markers for BC are available; carcinoembryonic antigen and cancer antigen 15.3 are the most used, but have low specificity and sensitivity [20]. Circulating tumour DNA and tumour cells seem very promising markers, however further studies are needed to validate them in routine clinical practice [21]. It would therefore be valuable if TPOAb could be confirmed as a blood BC prognostic marker.

Two studies evaluated 5-year outcomes in 142 [15] and 47 [16] BC women: Smyth *et al.* [15] reporting TPOAb- as a poor prognostic factor for disease-free survival (DFS) and overall survival (OS), and Fiore *et al.* [16] reporting 6.7% mortality in patients positive for anti-thyroid autoantibodies (TAb), mainly TPOAb+, compared with 46.9% in TAb negative patients. Farahati *et al.* evaluated 314 newly diagnosed BC patients and found no distant metastases among TPOAb+ patients compared with 6.6% among TPOAb- patients [17]. In contrast, Jiskra *et al.* followed 84 BC patients for 136 months (median), finding no impact of TPOAb on DFS or OS [19].

The aim of the present study was to clarify the impact of TPOAb on BC prognosis in a large, well powered patient cohort with long-term follow-up, according to the “REporting recommendations for tumour MARKer prognostic studies (REMARK)” guidelines [22]. The

“Taxotere as Adjuvant Chemotherapy Trial (TACT)” recruited 4162 women diagnosed with moderate-high risk early BC, evaluating whether sequential docetaxel (Taxotere) after anthracycline therapy would improve patient outcome compared with standard anthracycline chemotherapy: analyses were conducted at 62 months [23] and 97.5 months [24] follow-up, both showing no evidence of a difference between the two chemotherapy regimens. Of relevance, stored plasma was available in a significant number of these patients.

Furthermore, TPO is expressed in BC tissue [25], providing a possible mechanistic link: a thyroid/breast shared autoimmune response might target tumour cells and improve BC outcome. If TPOAb+ was confirmed as associated with a better BC outcome, new BC therapeutic approaches based on antigen-specific immunotherapies targeting TPO could be explored.

MATERIALS AND METHODS

Patients

The TACT study [23] was a multicentre, open-label, phase-III, randomised controlled trial of women aged >18 years diagnosed with operable early BC (pT1-3a, pN0-1, M0), with indication for adjuvant chemotherapy, including both lymph-node positive (node+) patients and lymph-node negative (node-) but high risk (e.g., tumour grade 3, hormonal-receptor expression negative, or lymphovascular invasion) patients.

Between February 2001 and June 2003, 4162 women were enrolled across 103 UK and one Belgian centres. All subjects underwent surgery, mastectomy or wide-local-excision (WLE), and were randomized (1:1 ratio) to the experimental regimen FEC-D (n=2073; fluorouracil, epirubicin, cyclophosphamide [FEC] followed by docetaxel) or centre’s choice of control chemotherapy, either FEC (n=1265) or E-CMF (n=824; epirubicin followed by CMF [cyclophosphamide, methotrexate, and fluorouracil]). Adjuvant radiotherapy was mandatory after WLE or used after mastectomy according to local guidelines. Endocrine treatments

(tamoxifen or aromatase-inhibitor monotherapy, tamoxifen followed by aromatase-inhibitor) were administered to patients with oestrogen receptor (ER) positive expression (ER+). Patients with human epidermal growth factor receptor-2 (HER2) positive expression (HER2+) were allowed to enter clinical trials assessing trastuzumab. All subjects have given their informed consent and the study protocol has been approved by the institute's committee on human research.

Laboratory measurements

Following a protocol amendment (November 2002), blood was taken for future translational research at the time of randomization, or at their next follow-up visit. Plasma samples were stored at -20°C for 6.5-13 years (range) at The Institute of Cancer Research (London, UK), and transferred to the Thyroid Research Group (Cardiff, UK) for TPOAb, thyrotropin (TSH) and free-thyroxine (FT4) analyses (October 2014) using an ADVIA Centaur automated immunoassay analyser (Bayer plc, UK) and Chemiluminescent Microparticle Immunoassay methods by the ARCHITECT® System (ABBOTT Laboratories, USA). According to the assay cut-off, TPOAb values were dichotomized as ≥ 6 kIU/L (positive: TPOAb+) versus < 6 kIU/L (negative: TPOAb-); TPOAb+ were also categorized into tertiles. FT4 and TSH normal ranges were 9.0–19.1 pmol/L and 0.30–4.40 mIU/L, respectively; they were also combined in a thyroid function status variable: euthyroid (FT4 and TSH within the normal ranges), hypothyroid (FT4 < 9.0 pmol/L and/or TSH > 4.40 mIU/L); hyperthyroid (FT4 > 19.1 pmol/L and/or TSH < 0.3 mIU/L).

Statistical analysis

According to TPOAb prevalence in age-matched females of general population [26,27], 20% of BC individuals were expected to be TPOAb+. Power calculations indicated 1158 and 1430 samples required to provide respectively 80% and 90% power to detect a 81% 5-year

DFS in TPOAb+ versus 73% in TPOAb- subjects (HR, 0.64; two-sided log-rank test with a 0.05 probability of a type I error), consistent with a 74.9% 5-year DFS rate in the whole TACT cohort [23].

Baseline characteristics, BC treatments and DFS-related characteristics were compared between TACT patients included or not in this study, and presented by dichotomized TPOAb and thyroid function status. Correlations between thyroid biomarkers were assessed using the Spearman rank method.

The primary outcome was to assess TPOAb prognostic significance in relation to DFS; secondary outcomes were TPOAb prognostic significance in relation to OS and time-to-recurrence (TTR), and thyroid function in relation to DFS, OS and TTR.

For DFS, OS and TTR, Kaplan-Meier curves were plotted and biomarkers compared with the log-rank test, and assessed firstly in a univariate Cox proportional hazards regression model stratified by centre's choice of control chemotherapy regimen and ER status, and subsequently included in a multivariable Cox model along with known BC prognostic factors: age, HER2 status, nodal involvement, tumour size and tumour grade. Additional variables, i.e. trial treatment (experimental versus control), type of surgery, trastuzumab use, radiotherapy and menopausal status, were included if, by stepwise selection ($P < 0.05$), shown to add value. TPOAb, TSH and FT4 were subsequently considered for inclusion if providing independent prognostic information. Interaction tests were used to explore differential effects within subgroups. HR with 95% CI were obtained, with $HR < 1$ indicating a better BC prognosis.

All patients with a biomarker value available were included in the analysis, as per an intention-to-treat analysis. All analyses were conducted using Stata version 13.1 (STATA CORP, TX) [23,24].

RESULTS

All available TACT plasma samples (N=2000) were analysed for thyroid biomarkers, and 1974 samples were considered for the statistical analyses (“analysis population”; **Supplemental Fig. 1**). The median (IQR; range) blood collection time was 15.5 (7.0-24.0; 0.5–57.2) months after surgery.

Supplemental Table 1 reports analysis population’s characteristics; the median (IQR; range) follow-up was 96.7 (87.4-106.3; 3.4-126.4) months. Overall 5-year estimates for DFS, OS and TTR were 79.5% (95% CI, 77.6-81.2), 87.4% (95% CI, 85.9-88.8) and 81.1% (95% CI, 79.3-82.8), respectively.

Distribution of TPOAb and thyroid function

TPOAb+ was detected in 406/1974 (20.6%) patients, distributed in the following tertiles: 137 (6.9%) 6-40 kIU/L (T1), 134 (6.7%) 41-238 kIU/L (T2), 135 (6.8%) 240-2000 kIU/L (T3). Baseline characteristics were largely comparable between TPOAb+ and TPOAb- patients (**Table 1**), apart from age, with TPOAb+ patients slightly older than TPOAb- patients (mean [SD] age, 50.2 [7.7] years versus 48.8 [8.5] years, respectively; P=0.005).

Plasma material was sufficient to determine FT4 and TSH values in 1974/1974 (100%) and 1971/1974 (99.8%) samples respectively. Among the 1974 patients, 1760 (89.2%) were euthyroid, 96 (4.9%) hypothyroid and 118 (6.0%) hyperthyroid; all 3 subgroups had similar baseline characteristics (**Table 1**), apart from age, with hypothyroid and hyperthyroid patients slightly older than euthyroid patients (mean [SD] age, respectively 50.5 [6.6] years and 50.7 [7.6] years, versus 48.9 [8.5] years; P=0.03).

As shown in **Supplemental Fig. 2**, FT4 and TSH were inversely correlated (Spearman rank, -0.23; P<0.001) and TPOAb was positively associated with TSH (Spearman rank, 0.24; P<0.001). The inverse correlation between TPOAb and FT4 was weak (Spearman rank, -0.04;

P=0.09). TPOAb+ cases were more prevalent among hypothyroid and hyperthyroid patients compared with the euthyroid group (73/96 [76.0%] hypothyroid; 45/118 [38.1%] hyperthyroid; 288/1760 [16.4%] euthyroid; P<0.001).

TPOAb and BC prognosis

The majority of DFS events were related to distant recurrence in both TPOAb+ and TPOAb- groups (**Supplemental Table 2**). There was no evidence of a difference in DFS between TPOAb+ and TPOAb- patients (unadjusted-HR: 0.97, 95% CI: 0.78-1.19, P=0.75, **Fig. 1A**; adjusted-HR: 1.00, 95% CI: 0.81-1.24, P=0.98, **Table 2**). Subgroup analyses showed no evidence of any significant interaction effects (**Fig. 2**). Similarly, there was no evidence of a difference by TPOAb status on OS (unadjusted-HR: 0.86, 95% CI: 0.66-1.11, P=0.24, **Fig. 1B**; adjusted-HR: 0.89, 95% CI: 0.69-1.14, P=0.35, not shown) and TTR (unadjusted-HR: 0.97, 95% CI: 0.78-1.21, P=0.80, **Fig. 1C**; adjusted-HR: 1.02, 95% CI: 0.81-1.27, P=0.89, not shown). TPOAb+ tertiles showed no evidence of a prognostic effect in both univariate (**Fig. 3**) and multivariable (data not shown) analyses for DFS, OS and TTR.

Two sensitivity analyses included 126 node+ patients not treated with radiotherapy, similar to Fiore *et al.* cohort [16], and 123 patients with blood taken before any adjuvant therapy. The median (IQR; range) time of blood collection after surgery was 12.4 (4.9-21.6; 0.7–47.2) months and 1.1 (0.9-1.4; 0.5-5.9) months, respectively. There was no evidence of a significant impact on DFS by TPOAb status in either of the two analyses, with unadjusted-HRs of 1.48 (95% CI, 0.68-3.25; P=0.32) and 0.83 (95% CI, 0.35-2.03; P=0.69) respectively.

Thyroid function and BC prognosis

There was no evidence of a significant difference for DFS, OS and TTR by thyroid function status in either univariate (**Fig. 4**) or multivariable (data not shown) analyses, and

when considering FT4 and TSH separately (DFS, **Supplemental Table 3**; OS and TTR, not shown).

DISCUSSION

In this large cohort of moderate-high risk early BC patients receiving adjuvant systemic treatments we found that neither the presence nor the titre of plasma TPOAb, assessed after BC diagnosis and measured with standard assays, had a substantial impact on long-term recurrence or mortality; similar findings were observed for thyroid status. These results confirm one previous finding [19], but contrast with two other studies [15,16]. We believe that our study is reliable, considering that our patient cohort is the largest to date, with one of the longest follow-ups, and focused on a well-defined BC population. Previous studies used smaller patient cohorts with shorter follow-ups [15,16,19], mixed different BC stages [19], or provided no information about BC stage [15], histological [15,19] and molecular subtypes [15,16,19], and adjuvant treatments received [15,19]; they may be susceptible to bias and random findings. In addition, the BC population analysed in this study is very similar to that of Fiore *et al.*, who recruited non-metastatic aggressive BC all treated with chemotherapy [16].

The long survival of our patient cohort could obscure a minor prognostic effect of TPOAb and/or thyroid function on BC, hypothetically detectable only among patients not suitable for standard treatments (e.g. medical contraindications) and targeted therapies (e.g. triple negative BC). This is possible but unlikely, since our exploratory analysis conducted among different BC subtypes confirmed our negative results. Furthermore, the multivariable analyses confirmed nodal status and tumour size as the two most important BC prognostic factors [28], proving that the cohort used was appropriate for the research question, and the model reasonably sensitive. Similarly, the better BC prognosis characterizing the intermediate age group (50-59 years) is consistent with the results of a recent large cohort study [29].

Our study cannot exclude a role of different TA parameters on BC prognosis, i.e. the presence of goitre [15] or incidental TA-related ¹⁸F-FDG PET/CT uptake [18]. Furthermore, differences in the alternative splicing of TPO in the breast as compared to the thyroid have been described [25], therefore this might also result in different TPO epitopes being targeted.

TPOAb prevalence in our cohort, similar to our *a priori* predicted value, reflects TPOAb prevalence among women of general population [26,30], increasing with age [26,31]. It remains possible that TPOAb+ rates are higher in the BC population, as our study was not designed to compare TPOAb prevalence among BC patients and the general population.

The principal limitations of the present study are the lack of clinical history for thyroid diseases or medications and that, similarly to previous studies [15,19], blood was mainly collected during/after adjuvant BC therapy. The first limitation might influence the prognostic role of thyroid function, but marginally of TPOAb, since they should exert an effect when either pre-existing, or appearing at a later time [32]; however, the evidence that thyroid function influences BC outcome is weak [6]. The finding of more cases of hyper- (6.0%) than hypothyroidism (4.9%) may reflect over-treatment with levothyroxine in some individuals.

Regarding BC adjuvant treatments, an increased risk of hypothyroidism after chemotherapy [33,34] or radiotherapy [35,36] for BC has been suggested in a few small studies, but not confirmed by others [37]. Tamoxifen can exert a modulation of thyroid function, mainly via an anti-thyroid effect [38,39] and the stress related to the surgical procedure itself has been suggested to cause immunomodulation [40]. However no clear large-scale effects of adjuvant treatments for BC, including trastuzumab, on thyroid function and immunity have been described, and our sensitivity analysis in a subgroup of 123 patients in whom blood was collected before BC adjuvant therapy showed no evidence of TPOAb prognostic ability, even if the wide 95% CI suggests a lack of statistical power.

To draw definitive conclusions, a prospective study collecting blood before cancer treatments would be ideal, but difficult to realise because of the large patient number required,

as shown by our *a priori* power calculation. Furthermore, this study analysed moderate-high risk early BC only. BC is a heterogeneous disease, with many subtypes characterised by different clinical behaviour and prognosis; it could be possible that TPOAb and/or thyroid function affect the prognosis of certain specific BC subtypes and stages only, therefore they should be all investigated separately, with a much higher total patient numbers required to reach significant and definitive results.

In conclusion, the present study is to our knowledge the largest currently available investigating the impact of blood TPOAb and thyroid function on BC prognosis, providing a detailed description of the BC population analysed, and therefore representing a key-work to clarify this debate over decades. We found that TPOAb and thyroid function, both measured with standard assays and after BC diagnosis, appear not to influence substantially the long-term recurrence and mortality of moderate-high risk early BC in the modern era. Major confounding in this conclusion due to BC treatments seems unlikely. Future studies might explore different BC stages and/or specific subtypes, also searching for non-conventional or breast-specific immune responses to particular TPO epitopes, to determine whether aspects of TA other than standard TPOAb and thyroid function may be relevant to BC outcome.

274 **ACKNOWLEDGMENTS**

275 This study was supported by the Tenovus Cancer Care, Cardiff, UK (2013 Tenovus Innovation
276 Grant number TIG2014-28).

277 Authors thank all the patients who participated in this study and all the investigators involved
278 in the TACT trial (funded by Cancer Research UK; ICR-CTSU also receives core funding from
279 Cancer Research UK).

280 Authors thank Dr. C Evans and her team (Department of Medical Biochemistry, School of
281 Medicine, Cardiff University, UK) for running the laboratory assays.

REFERENCES

1. Hardefeldt PJ, Eslick GD, Edirimanne S: Benign thyroid disease is associated with breast cancer: a meta-analysis. *Breast Cancer Res Treat* 2012;133:1169-77.
2. Prinzi N, Baldini E, Sorrenti S, De Vito C, Tuccilli C, Catania A, Carbotta S, Mocini R, Cocco C, Nesca A, Bianchini M, De Antoni E, D'Armiento M, Ulisse S: Prevalence of breast cancer in thyroid diseases: results of a cross-sectional study of 3,921 patients. *Breast Cancer Res Treat* 2014;144:683-8.
3. Sarlis NJ, Gourgiotis L, Pucino F, Tolis GJ: Lack of association between Hashimoto thyroiditis and breast cancer: a quantitative research synthesis. *Hormones (Athens)* 2002;1:35-41.
4. Angelousi AG, Anagnostou VK, Stamatakis MK, Georgiopoulos GA, Kontzoglou KC: Mechanisms in endocrinology: primary HT and risk for breast cancer: a systematic review and meta-analysis. *Eur J Endocrinol* 2012;166:373-81.
5. Moeller LC, Fuhrer D, Moeller LC, Fuhrer D: Thyroid hormone, thyroid hormone receptors, and cancer: a clinical perspective. *Endocrine-Related Cancer* 2013;20:R19-29.
6. Smyth PP: The thyroid and breast cancer. *Curr Opin Endocrinol Diabetes Obes* 2016;23:389-93.
7. Mittra I, Hayward JL: Hypothalamic-pituitary-thyroid axis in breast cancer. *Lancet* 1974;1:885-9.
8. Kuijpers JL, Nyklicek I, Louwman MW, Weetman TA, Pop VJ, Coebergh JW: Hypothyroidism might be related to breast cancer in post-menopausal women. *Thyroid* 2005;15:1253-9.
9. Cristofanilli M, Yamamura Y, Kau SW, Bevers T, Strom S, Patangan M, Hsu L, Krishnamurthy S, Theriault RL, Hortobagyi GN: Thyroid hormone and breast carcinoma.

307 Primary hypothyroidism is associated with a reduced incidence of primary breast carcinoma.
308 Cancer 2005;103:1122-8.

309 10. Tosovic A, Becker C, Bondeson AG, Bondeson L, Ericsson UB, Malm J,
310 Manjer J: Prospectively measured thyroid hormones and thyroid peroxidase antibodies in
311 relation to breast cancer risk. Int J Cancer 2012;131:2126-33.

312 11. Sogaard M, Farkas DK, Ehrenstein V, Jorgensen JO, Dekkers OM, Sorensen
313 HT: Hypothyroidism and hyperthyroidism and breast cancer risk: a nationwide cohort study.
314 Eur J Endocrinol 2016;174:409-14.

315 12. Hedley AJ, Jones SJ, Spiegelhalter DJ, Clements P, Bewsher PD, Simpson JG,
316 Weir RD: Breast cancer in thyroid disease: fact or fallacy? Lancet 1981;1:131-3.

317 13. Hellevik AI, Asvold BO, Bjoro T, Romundstad PR, Nilsen TI, Vatten LJ:
318 Thyroid function and cancer risk: a prospective population study. Cancer Epidemiol
319 Biomarkers Prev 2009;18:570-4.

320 14. Giani C, Fierabracci P, Bonacci R, Gigliotti A, Campani D, De Negri F,
321 Cecchetti D, Martino E, Pinchera A: Relationship between breast cancer and thyroid disease:
322 relevance of autoimmune thyroid disorders in breast malignancy. J Clin Endocrinol Metab
323 1996;81:990-4.

324 15. Smyth PP, Shering SG, Kilbane MT, Murray MJ, McDermott EW, Smith DF,
325 O'Higgins NJ: Serum thyroid peroxidase autoantibodies, thyroid volume, and outcome in
326 breast carcinoma. J Clin Endocrinol Metab 1998;83:2711-6.

327 16. Fiore E, Giustarini E, Mammoli C, Fragomeni F, Campani D, Muller I, Pinchera
328 A, Giani C: Favorable predictive value of thyroid autoimmunity in high aggressive breast
329 cancer. J Endocrinol Invest 2007;30:734-8.

330 17. Farahati J, Roggenbuck D, Gilman E, Schutte M, Jagminaitė E, Seyed Zakavi
331 R, Loning T, Heissen E: Anti-thyroid peroxidase antibodies are associated with the absence of

332 distant metastases in patients with newly diagnosed breast cancer. Clin Chem Lab Med
333 2012;50:709-14.

334 18. Kim SS, Kim IJ, Kim SJ, Lee JY, Bae YT, Jeon YK, Kim BH, Kim YK:
335 Incidental diffuse thyroid 18F-FDG uptake related to autoimmune thyroiditis may be a
336 favorable prognostic factor in advanced breast cancer. J Nucl Med 2012;53:1855-62.

337 19. Jiskra J, Barkmanova J, Limanova Z, Lanska V, Smutek D, Potlukova E,
338 Antosova M: Thyroid autoimmunity occurs more frequently in women with breast cancer
339 compared to women with colorectal cancer and controls but it has no impact on relapse-free
340 and overall survival. Oncol Rep 2007;18:1603-11.

341 20. Cheung KL, Graves CR, Robertson JF: Tumour marker measurements in the
342 diagnosis and monitoring of breast cancer. Cancer Treat Rev 2000;26:91-102.

343 21. Ignatiadis M, Lee M, Jeffrey SS: Circulating Tumor Cells and Circulating
344 Tumor DNA: Challenges and Opportunities on the Path to Clinical Utility. Clin Cancer Res
345 2015;21:4786-800.

346 22. McShane LM, Altman DG, Sauerbrei W, Taube SE, Gion M, Clark GM,
347 Statistics Subcommittee of the NCIEWGoCD: REporting recommendations for tumor
348 MARKer prognostic studies (REMARK). Nat Clin Pract Oncol 2005;2:416-22.

349 23. Ellis P, Barrett-Lee P, Johnson L, Cameron D, Wardley A, O'Reilly S, Verrill
350 M, Smith I, Yarnold J, Coleman R, Earl H, Canney P, Twelves C, Poole C, Bloomfield D,
351 Hopwood P, Johnston S, Dowsett M, Bartlett JM, Ellis I, Peckitt C, Hall E, Bliss JM, Group
352 TTM, Trialists T: Sequential docetaxel as adjuvant chemotherapy for early breast cancer
353 (TACT): an open-label, phase III, randomised controlled trial. Lancet 2009;373:1681-92.

354 24. Bliss JM, Ellis P, Kilburn L, Bartlett J, Bloomfield D, Cameron D, Canney P,
355 Coleman RE, Dowsett M, Earl H, Verrill M, Wardley A, Yarnold J, Ahern R, Atkins N,
356 Fletcher M, McLinden M, Barrett-Lee P: Mature analysis of UK Taxotere as Adjuvant

357 Chemotherapy (TACT) trial (CRUK 01/001); effects of treatment and characterisation of
358 patterns of breast cancer relapse. *Cancer Res Suppl.* 2012;72:Abstract P1-13-03.

359 25. Muller I, Giani C, Zhang L, Grennan-Jones FA, Fiore E, Belardi V, Rosellini
360 V, Funel N, Campani D, Giustarini E, Lewis MD, Bakhsh AD, Roncella M, Ghilli M, Vitti P,
361 Dayan CM, Ludgate ME: Does thyroid peroxidase provide an antigenic link between thyroid
362 autoimmunity and breast cancer? *Int J Cancer* 2014;134:1706-14.

363 26. Vanderpump MP, Tunbridge WM, French JM, Appleton D, Bates D, Clark F,
364 Grimley Evans J, Hasan DM, Rodgers H, Tunbridge F, et al.: The incidence of thyroid
365 disorders in the community: a twenty-year follow-up of the Wickham Survey. *Clin*
366 *Endocrinol (Oxf)* 1995;43:55-68.

367 27. Bulow Pedersen I, Knudsen N, Carle A, Vejbjerg P, Jorgensen T, Perrild H,
368 Ovesen L, Banke Rasmussen L, Laurberg P: A cautious iodization program bringing iodine
369 intake to a low recommended level is associated with an increase in the prevalence of thyroid
370 autoantibodies in the population. *Clin Endocrinol (Oxf)* 2011;75:120-6.

371 28. Fitzgibbons PL, Page DL, Weaver D, Thor AD, Allred DC, Clark GM, Ruby
372 SG, O'Malley F, Simpson JF, Connolly JL, Hayes DF, Edge SB, Lichter A, Schnitt SJ:
373 Prognostic factors in breast cancer. College of American Pathologists Consensus Statement
374 1999. *Arch Pathol Lab Med* 2000;124:966-78.

375 29. Brandt J, Garne JP, Tengrup I, Manjer J: Age at diagnosis in relation to survival
376 following breast cancer: a cohort study. *World J Surg Oncol* 2015;13:33.

377 30. Hollowell JG, Staehling NW, Flanders WD, Hannon WH, Gunter EW, Spencer
378 CA, Braverman LE: Serum TSH, T(4), and thyroid antibodies in the United States population
379 (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III). *J Clin*
380 *Endocrinol Metab* 2002;87:489-99.

- 381 31. Tunbridge WM, Brewis M, French JM, Appleton D, Bird T, Clark F, Evered
382 DC, Evans JG, Hall R, Smith P, Stephenson J, Young E: Natural history of autoimmune
383 thyroiditis. *Br Med J (Clin Res Ed)* 1981;282:258-62.
- 384 32. Franzke A, Peest D, Probst-Kepper M, Buer J, Kirchner GI, Brabant G,
385 Kirchner H, Ganser A, Atzpodien J: Autoimmunity resulting from cytokine treatment predicts
386 long-term survival in patients with metastatic renal cell cancer. *J Clin Oncol* 1999;17:529-33.
- 387 33. Kumar N, Allen KA, Riccardi D, Bercu BB, Cantor A, Minton S, Balducci L,
388 Jacobsen PB: Fatigue, weight gain, lethargy and amenorrhea in breast cancer patients on
389 chemotherapy: is subclinical hypothyroidism the culprit? *Breast Cancer Res Treat*
390 2004;83:149-59.
- 391 34. de Groot S, Janssen LGM, Charehbili A, Dijkgraaf EM, Smit VTHBM, Kessels
392 LW, van Bochove A, van Laarhoven HWM, Meershoek-Klein Kranenbarg E, van Leeuwen-
393 Stok AE, van de Velde CJH, Putter H, Nortier JWR, van der Hoeven JJM, Pijl H, Kroep JR:
394 Thyroid function alters during neoadjuvant chemotherapy in breast cancer patients: results
395 from the NEOZOTAC trial (BOOG 2010-01). *Breast Cancer Research and Treatment*
396 2015;149:461-466.
- 397 35. Bruning P, Bonfrer J, De Jong-Bakker M, Nooyen W, Burgers M: Primary
398 hypothyroidism in breast cancer patients with irradiated supraclavicular lymph nodes. *Br J*
399 *Cancer* 1985;51:659-63.
- 400 36. Cutuli B, Quentin P, Rodier JF, Barakat P, Grob JC: Severe hypothyroidism
401 after chemotherapy and locoregional irradiation for breast cancer. *Radiother Oncol*
402 2000;57:103-5.
- 403 37. Smith GL, Smith BD, Giordano SH, Shih YC, Woodward WA, Strom EA,
404 Perkins GH, Tereffe W, Yu TK, Buchholz TA: Risk of hypothyroidism in older breast cancer
405 patients treated with radiation. *Cancer* 2008;112:1371-9.

- 406 38. Anker GB, Lonning PE, Aakvaag A, Lien EA: Thyroid function in
407 postmenopausal breast cancer patients treated with tamoxifen. Scand J Clin Lab Invest
408 1998;58:103-7.
- 409 39. Zidan J, Rubenstein W: Effect of adjuvant tamoxifen therapy on thyroid
410 function in postmenopausal women with breast cancer. Oncology 1999;56:43-5.
- 411 40. Boomsma MF, Garssen B, Slot E, Berbee M, Berkhof J, Meezenbroek Ede J,
412 Slieker W, Visser A, Meijer S, Beelen RH: Breast cancer surgery-induced immunomodulation.
413 J Surg Oncol 2010;102:640-8.
- 414

416 Table 1: Baseline characteristics and treatments for breast cancer by autoantibodies to thyroid peroxidase (TPOAb) and thyroid function status

	TPOAb- N = 1568	TPOAb+ N = 406	P value	Hypothyroid N = 96	Euthyroid N = 1760	Hyperthyroid N = 118	P value
Age (years): mean (SD)	48.8 (8.5)	50.2 (7.7)	0.005 ^a	50.5 (6.6)	48.9 (8.5)	50.7 (7.6)	0.03 ^d
Age group (years): n (%)			0.08 ^b				0.62 ^b
<40	257 (16.4)	49 (12.1)		8 (8.3)	287 (16.3)	11 (9.3)	
40-49	575 (36.7)	151 (37.2)		36 (37.5)	647 (36.8)	43 (36.4)	
50-59	590 (37.6)	167 (41.1)		45 (46.9)	657 (37.3)	55 (46.6)	
≥60	146 (9.3)	39 (9.6)		7 (7.3)	169 (9.6)	9 (7.6)	
Nodal status: n (%)			0.62 ^b				0.61 ^b
Node negative	314 (20.0)	93 (22.9)		18 (18.8)	367 (20.9)	22 (18.6)	
1-3 positive nodes	719 (45.9)	171 (42.1)		33 (34.4)	808 (45.9)	49 (41.5)	
≥4 positive nodes	535 (34.1)	142 (35.0)		45 (46.9)	585 (33.2)	47 (39.8)	
Tumour grade: n (%)			0.74 ^b				0.72 ^b
Grade 1	77 (4.9)	23 (5.7)		4 (4.2)	88 (5.0)	8 (6.8)	
Grade 2	603 (38.5)	155 (38.2)		35 (36.5)	681 (38.7)	42 (35.6)	
Grade 3	883 (56.3)	228 (56.2)		57 (59.4)	986 (56.0)	68 (57.6)	
Unknown	5 (0.3)	0 (0.0)		0 (0.0)	5 (0.3)	0 (0.0)	
Tumour size (cm): n (%)			0.59 ^b				0.38 ^b
≤2	578 (36.9)	147 (36.2)		25 (26.0)	659 (37.4)	41 (34.8)	
>2 and ≤5	857 (54.7)	220 (54.2)		61 (63.5)	952 (54.1)	64 (54.2)	
>5	132 (8.4)	39 (9.6)		10 (10.4)	148 (8.4)	13 (11.0)	
Unknown	1 (0.1)	0 (0.0)		0 (0.0)	1 (0.1)	0 (0.0)	
ER & HER2 status: n (%)			0.85 ^c (ER) 0.45 ^c (HER2)				0.62 ^c (ER) 0.84 ^c (HER2)
ER+	1107 (70.6)	289 (71.2)		69 (71.9)	1248 (70.9)	79 (67.0)	
& HER2+	198 (12.6)	49 (12.1)		13 (13.5)	220 (12.5)	14 (11.9)	
& HER2-	772 (49.2)	201 (49.5)		46 (47.9)	873 (49.6)	54 (45.8)	
& HER2 unknown	137 (8.7)	39 (9.6)		10 (10.4)	155 (8.8)	11 (9.3)	
ER-	461 (29.4)	117 (28.8)		27 (28.1)	512 (29.1)	39 (33.1)	
& HER2+	118 (7.5)	43 (10.6)		8 (8.3)	141 (8.0)	12 (10.2)	
& HER2-	289 (18.4)	61 (15.0)		15 (15.6)	313 (17.8)	22 (18.6)	
& HER2 unknown	54 (3.4)	13 (3.2)		4 (4.2)	58 (3.3)	5 (4.2)	
Molecular subgroup: n (%)			0.40 ^c				0.94 ^c
ER+/HER2- ¹	784 (50.0)	203 (50.0)		47 (49.0)	885 (50.3)	55 (46.6)	
HER2+	316 (20.2)	92 (22.7)		21 (21.9)	361 (20.5)	26 (22.0)	
Triple negative	277 (17.7)	59 (14.5)		14 (14.6)	301 (17.1)	21 (17.8)	

	TPOAb- N = 1568	TPOAb+ N = 406	P value	Hypothyroid N = 96	Euthyroid N = 1760	Hyperthyroid N = 118	P value
Type of surgery and radiotherapy use: n (%)							
Mastectomy	854 (54.5)	225 (55.4)	0.74 ^c (surgery) 0.61 ^c (radiotherapy)	53 (55.2)	962 (54.7)	64 (54.2)	0.99 ^c (surgery) 0.33 ^c (radiotherapy)
with radiotherapy [^]	688 (80.6)	177 (78.7)		47 (88.7)	772 (80.2)	46 (71.9)	
Wide local excision	714 (45.5)	181 (44.6)		43 (44.8)	798 (45.3)	54 (45.8)	
with radiotherapy [#]	704 (98.6)	176 (97.2)		41 (95.3)	787 (98.6)	52 (96.3)	
Endocrine treatment in ER+ patients: n (%)*							
Tamoxifen monotherapy	696 (62.9)	167 (57.8)	0.13 ^c	43 (62.3)	772 (61.9)	48 (60.8)	0.09 ^c
Tamoxifen followed by AI	354 (32.0)	100 (34.6)		20 (29.0)	409 (32.8)	25 (31.7)	
AI monotherapy	46 (4.2)	15 (5.2)		6 (8.7)	53 (4.3)	2 (2.5)	
No endocrine treatment/unknown	11 (1.0)	7 (2.4)		0 (0.0)	14 (1.1)	4 (5.1)	
Trastuzumab in HER2+ patients: n (%)**							
Yes	40 (12.7)	8 (8.7)	0.36 ^c	1 (4.8)	44 (12.2)	3 (11.5)	0.71 ^c
No/Not known	276 (87.3)	84 (91.3)		20 (95.2)	317 (87.8)	23 (88.5)	
Chemotherapy: n (%)							
Control (FEC)	498 (31.8)	128 (31.5)	0.52 ^c	27 (28.1)	568 (32.3)	31 (26.3)	0.90 ^c
Control (E-CMF)	271 (17.3)	61 (15.0)		16 (16.7)	301 (17.1)	15 (12.7)	
FEC-D	799 (51.0)	217 (53.4)		53 (55.2)	891 (50.6)	52 (44.1)	

¹ includes ER-, PgR+, HER2-

[^] denominators calculated using patients treated with mastectomy

[#] denominators calculated using patients treated with wide local excision

^{*} denominators calculated using ER+ patients

^{**} denominators calculated using HER2+ patients

^a t-test

^b trend test; note “unknowns” excluded from the test

^c Fisher’s exact test

^d ANOVA

AI, aromatase-inhibitors; ER+, positive estrogen receptor (ER); ER-, negative ER; E-CMF, epirubicin 100 mg/m² for 4 cycles followed by CMF (cyclophosphamide 600 mg/m², methotrexate 40 mg/m² and fluorouracil 600 mg/m²) for 4 cycles; FEC, fluorouracil 600 mg/m², epirubicin 60 mg/m² and cyclophosphamide 600 mg/m² for 8 cycles; FEC-D, FEC for 4 cycles followed by docetaxel 100 mg/m² for 4 cycles; HER2+, positive human epidermal growth factor receptor-2 (HER2); HER2-, negative HER2; PgR+, positive progesterone receptor (PgR); SD, standard deviation; TPOAb+, positive TPOAb; TPOAb-, negative TPOAb; triple negative, negative HER2, ER and PgR.

Table 2: Multivariable analysis for disease-free survival by dichotomized autoantibodies to thyroid peroxidase (TPOAb)


		HR	95% CI	P value
TPOAb status	negative (n=1568)	1.00	-	-
	positive (n=406)	1.00	0.81-1.24	0.98
Nodal status	positive (n=1567)	1.00	-	-
	negative (n=407)	0.49	0.37-0.64	< 0.001
HER2 status	negative (n=1323)	1.00	-	-
	positive (n=408)	1.19	0.97-1.46	0.09
	unknown (n=243)	0.93	0.71-1.23	0.63
Age group (years)	<40 (n=306)	1.00	-	-
	40-49 (n=726)	0.78	0.61-1.00	0.05
	50-59 (n=757)	0.75	0.59-0.96	0.02
	≥60 (n=185)	0.95	0.69-1.31	0.76
Tumour grade	Grade 1 (n=100)	1.00	-	-
	Grade 2 (n=758)	1.15	0.74-1.78	0.55
	Grade 3 (n=1111)	1.39	0.89-2.17	0.14
	unknown (n=5)	0.77	0.10-5.75	0.80
Tumour size (cm) *	≤2 (n=725)	1.00	-	-
	>2 and ≤5 (n=1077)	1.37	1.12-1.66	0.002
	>5 (n=171)	1.88	1.41-2.52	< 0.001
Type of surgery	Mastectomy (n=1079)	1.00	-	-
	WLE (n=895)	0.79	0.66-0.95	0.01

HER2, human epidermal growth factor receptor-2; HR, hazard ratio (HR <1 indicates a favorable breast cancer outcome); WLE, wide local excision; 95% CI, 95% confidence interval.

* The patient with unknown tumour size (n=1) has not been considered for this analysis.


FIGURES

Fig. 1: Univariate analyses by dichotomized autoantibodies to thyroid peroxidase (TPOAb)




Kaplan-Meier curves relative to breast cancer (BC) outcome (median follow-up 96.7 months) in patients positive (≥ 6 kIU/L) and negative (< 6 kIU/L) for TPOAb. HR, hazard ratio (HR < 1 indicates a favorable BC outcome); 95% CI, 95% confidence interval. Panel A: disease-free survival (DFS). Panel B: overall survival (OS). Panel C: time to recurrence (TTR).

Fig. 2: Exploratory subgroup analyses for disease-free survival by dichotomized autoantibodies to thyroid peroxidase (TPOAb)




ER+, positive estrogen receptor (ER); ER-, negative ER; HER2+, positive human epidermal growth factor receptor-2 (HER2); HER2-, negative HER2; NK, not known; N-, lymph-node negative; N1-3+, 1-3 lymph-nodes positive, N4+, 4 or more lymph-nodes positive; TPOAb+, positive TPOAb; TPOAb-, negative TPOAb; triple negative, negative HER2, ER and progesterone receptor; WLE, wide local excision; 95% CI, 95% confidence interval.

Fig. 3: Univariate analyses by autoantibodies to thyroid peroxidase (TPOAb) categorized into tertiles



Kaplan-Meier curves relative to breast cancer (BC) outcome (median follow-up 96.7 months) in patients negative (<6 kIU/L) and positive for TPOAb categorized into tertiles: 6-40 kIU/L (T1), 41-238 kIU/L (T2), 240-2000 kIU/L (T3). HR, hazard ratio (HR <1 indicates a favorable BC outcome); 95% CI, 95% confidence interval. Panel A: disease-free survival (DFS). Panel B: overall survival (OS). Panel C: time to recurrence (TTR).

Fig. 4: Univariate analyses by thyroid function status



Kaplan-Meier curves relative to breast cancer (BC) outcome (median follow-up 96.7 months) according to thyroid function status. Euthyroid, free-thyroxine (FT4) 9.0–19.1 pmol/L and thyrotropin (TSH) 0.30–4.40 mIU/L; hyperthyroid, FT4 >19.1 pmol/L and/or TSH <0.3 mIU/L; hypothyroid, FT4 <9.0 pmol/L and/or TSH >4.40 mIU/L. HR, hazard ratio (HR <1 indicates a favorable BC outcome); 95% CI, 95% confidence interval. Panel A: disease-free survival (DFS). Panel B: overall survival (OS). Panel C: time to recurrence (TTR).

477 **SUPPLEMENTAL MATERIAL**

478

479 **Supplemental Table 1: Baseline characteristics, treatments for breast cancer and disease-free**

480 **survival (DFS) related characteristics**

481

	Analysis population N = 1974	Not included patients N = 2188	All TACT trial patients N = 4162
Age (years), mean (SD)	49.1 (8.4)	48.2 (8.6)	48.6 (8.5)
Age group (years), n (%)			
<40	306 (15.5)	412 (18.8)	718 (17.3)
40-49	726 (36.8)	841 (38.4)	1567 (37.7)
50-59	757 (38.4)	730 (33.4)	1487 (35.7)
≥60	185 (9.4)	205 (9.4)	390 (9.4)
Nodal status, n (%)			
Node negative	407 (20.6)	428 (19.6)	835 (20.1)
1-3 positive nodes	890 (45.1)	949 (43.4)	1839 (44.2)
≥4 positive nodes	677 (34.3)	811 (37.1)	1488 (35.8)
Tumor grade, n (%)			
Grade 1	100 (5.1)	129 (5.9)	229 (5.5)
Grade 2	758 (38.4)	778 (35.6)	1536 (36.9)
Grade 3	1111 (56.3)	1271 (58.1)	2382 (57.2)
Unknown	5 (0.3)	10 (0.5)	15 (0.4)
Tumor size (cm), n (%)			
≤2	725 (36.7)	711 (32.5)	1436 (34.5)
>2 and ≤5	1077 (54.6)	1253 (57.3)	2330 (56.0)
>5	171 (8.7)	221 (10.1)	392 (9.4)
Unknown	1 (0.1)	3 (0.1)	4 (0.1)
ER & HER2 status, n (%)			
ER+	1396 (70.7)	1479 (67.6)	2875 (69.1)
& HER2+	247 (12.5)	247 (11.3)	494 (11.9)
& HER2-	973 (49.3)	990 (45.2)	1963 (47.2)
& HER2 unknown	176 (8.9)	242 (11.1)	418 (10.0)
ER-	578 (29.3)	709 (32.4)	1287 (30.9)
& HER2+	161 (8.2)	194 (8.9)	355 (8.5)
& HER2-	350 (17.7)	411 (18.8)	761 (18.3)
& HER2 unknown	67 (3.4)	104 (4.8)	171 (4.1)
Molecular subgroup, n (%)			
ER+/HER2-*	987 (50.0)	1014 (46.3)	2001 (48.1)
HER2+	408 (20.7)	441 (20.2)	849 (20.4)
Triple negative	336 (17.0)	387 (17.7)	723 (17.4)
Type of surgery and radiotherapy, n (%)			
Mastectomy	1079 (54.7)	1186 (54.2)	2265 (54.4)
with radiotherapy	865 (43.8)	949 (43.4)	1814 (43.6)
breast	159 (8.1)	254 (11.6)	413 (9.9)
chest wall	709 (35.9)	693 (31.7)	1402 (33.7)
supraclavicular fossa	480 (24.3)	500 (22.9)	980 (23.5)
axilla	85 (4.3)	103 (4.7)	188 (4.5)
Wide local excision	895 (45.3)	1002 (45.8)	1897 (45.6)
with radiotherapy	880 (44.6)	961 (43.9)	1841 (44.2)
breast	856 (43.4)	921 (42.1)	1777 (42.7)
chest wall	31 (1.6)	47 (2.1)	78 (1.9)
supraclavicular fossa	291 (14.7)	283 (12.9)	574 (13.8)
axilla	103 (5.2)	70 (3.2)	173 (4.2)
Endocrine treatment in ER+ patients, n (%)			
Tamoxifen monotherapy	863 (61.8)	927 (62.7)	1790 (62.3)

	Analysis population N = 1974	Not included patients N = 2188	All TACT trial patients N = 4162
Tamoxifen followed by AI	454 (32.5)	439 (29.7)	893 (31.1)
AI monotherapy	61 (4.4)	76 (5.1)	137 (4.8)
No endocrine treatment/unknown	18 (1.3)	37 (2.5)	55 (1.9)
Trastuzumab in HER2+ patients, n (%)			
Yes	48 (11.8)	28 (6.4)	76 (9.0)
No/Not known	360 (88.2)	413 (93.7)	773 (91.0)
Chemotherapy, n (%)			
Control (FEC)	626 (31.7)	639 (29.2)	1265 (30.4)
Control (E-CMF)	332 (16.8)	492 (22.5)	824 (19.8)
FEC-D	1016 (51.5)	1057 (48.3)	2073 (49.5)
Number of patients with event contributing to DFS analysis	551 (27.9)	778 (35.6)	1329 (31.9)
Local recurrence	76 (3.8)	107 (4.9)	183 (4.4)
Distant recurrence	405 (20.5)	572 (26.1)	977 (23.5)
New breast disease	43 (2.2)	44 (2.0)	91 (2.2)
Death from other cause (no recurrence)	27 (1.4)	51 (2.3)	78 (1.9)
Distant relapse ever reported	462 (23.4)	655 (29.9)	1117 (26.8)
New breast disease ever reported	57 (2.9)	67 (3.1)	124 (3.0)
All non-breast cancer second primary	52 (2.6)	54 (2.5)	106 (2.5)
All deaths	397 (20.1)	620 (28.3)	1017 (24.4)
Breast cancer	369 (18.7)	568 (26.0)	937 (22.5)
Death from other causes	28 (1.4)	52 (2.4)	80 (1.9)
Cancer (non-breast)	15 (0.8)	21 (1.0)	36 (0.9)
Treatment toxicity	0	5 (0.2)	5 (0.1)
Other	13 (0.7)	26 (1.2)	39 (0.9)

* includes ER-, PgR+, HER2-

AI, aromatase-inhibitors; ER+, positive estrogen receptor (ER); ER-, negative ER; E-CMF, epirubicin 100 mg/m² for 4 cycles followed by CMF (cyclophosphamide 600 mg/m², methotrexate 40 mg/m² and fluorouracil 600 mg/m²) for 4 cycles; FEC, fluorouracil 600 mg/m², epirubicin 60 mg/m² and cyclophosphamide 600 mg/m² for 8 cycles; FEC-D, FEC for 4 cycles followed by docetaxel 100 mg/m² for 4 cycles; HER2+, positive human epidermal growth factor receptor-2 (HER2); HER2-, negative HER2; PgR+, positive progesterone receptor (PgR); SD, standard deviation; TACT, "Taxotere as adjuvant chemotherapy trial"; TPOAb, autoantibodies to thyroid peroxidase.

Supplemental Table 2: Events contributing to disease-free survival (DFS) and numbers of deaths by dichotomized TPOAb status

	TPOAb- (N = 1568) n (%)	TPOAb+ (N = 406) n (%)
Number of patients with event contributing to DFS analysis	442 (28.2)	109 (26.8)
Local recurrence	59 (3.8)	17 (4.2)
Distant recurrence	327 (20.9)	78 (19.2)
New breast disease	33 (2.1)	10 (2.5)
Death from other cause (no recurrence)	23 (1.5)	4 (1.0)
All deaths	325 (20.7)	72 (17.7)
Breast cancer	301 (19.2)	68 (16.7)
Death from other causes (without distant recurrence)	24 (1.5)	4 (1.0)
Cancer (non-breast)	14 (0.9)	1 (0.2)
Treatment toxicity	0 (0.0)	0 (0.0)
Other	9 (0.6)	3 (0.7)
Vascular (cardiac)	1 (0.1)	1 (0.2)
Vascular (cerebral)	1 (0.1)	0 (0.0)
Vascular (thromboembolic)	0 (0.0)	0 (0.0)
Respiratory	0 (0.0)	0 (0.0)
Accident, suicide, alcoholism	5 (0.3)	0 (0.0)
Infection (not treatment related)	0 (0.0)	1 (0.2)
Gastrointestinal bleed	0 (0.0)	0 (0.0)
Chronic liver disease	1 (0.1)	0 (0.0)
Unknown	2 (0.1)	1 (0.2)

TPOAb+, positive autoantibodies to thyroid peroxidase (TPOAb); TPOAb-, negative TPOAb.


499 **Supplemental Table 3: Univariate analyses for disease-free survival by FT4 and TSH**

Variable		Unadjusted HR	95% CI	P value
FT4	Continuous	1.00	0.96-1.04	0.91
	<9.0 pmol/L (Hypothyroid; n=13)	1.61	0.67-3.88	0.29
	9.0–19.1 pmol/L (Euthyroid; n=1917)	1.00	-	-
	>19.1 pmol/L (Hyperthyroid; n=44)	1.08	0.62-1.87	0.79
TSH*	Continuous	1.03	0.94-1.13	0.48
	>4.40 mIU/L (Hypothyroid; n=94)	1.08	0.73-1.59	0.71
	0.3–4.40 mIU/L (Euthyroid; n=1781)	1.00	-	-
	<0.3 mIU/L (Hyperthyroid; n=96)	1.19	0.82-1.72	0.36

500 FT4, free-thyroxine; HR, hazard ratio (HR <1 indicates a favorable breast cancer outcome); TSH,
501 thyrotropin; 95% CI, 95% confidence interval.


502
503 *TSH value was available in 1971/1974 (99.8%) samples
504

Supplemental Fig. 1: Flowchart regarding sample availability and data handling



FT4, free-thyroxine; TPOAb, autoantibodies to thyroid peroxidase; TSH, thyrotropin.

509 **Supplemental Fig. 2: Correlation between thyroid markers**



510

511 FT4, free-thyroxine; TPOAb, autoantibodies to thyroid peroxidase; TSH, thyrotropin. Panel A:

512 inverse correlation between TSH (logarithmic scale) and FT4. Panel B: positive correlation

513 between TPOAb and TSH (logarithmic scale). Panel C: inverse correlation between TPOAb
514 and FT4.